arXiv:1502.06131v2 [math.CO] 18 Feb 2016

UNIMODULAR BINARY HIERARCHICAL MODELS

DANIEL IRVING BERNSTEIN AND SETH SULLIVANT

Abstract. Associated to each simplicial complex is a binary hierarchical model. We classify the simplicial complexes that yield unimodular binary hierarchical models. Our main theorem provides both a construction of all unimodular binary hierarchical models, together with a characterization in terms of excluded minors, where our definition of a minor allows the taking of links and induced complexes. A key tool in the proof is the lemma that the class of unimodular binary hierarchical models is closed under the Alexander duality operation on simplicial complexes.

1. Introduction

Associated to a simplicial complex $\mathcal{C}$ with ground set $[m]$ and an integer vector $\mathbf{d} \in \mathbb{Z}^m$ is an integral matrix $\mathcal{A}_{\mathcal{C},\mathbf{d}}$. The hierarchical model associated to $\mathcal{C}, \mathbf{d}$ is a log-linear model (i.e. discrete exponential family) whose design matrix is the matrix $\mathcal{A}_{\mathcal{C},\mathbf{d}}$. The Zariski closure of a hierarchical model is a (not necessarily normal) toric variety. As for all log-linear models, important relevant problems are to study properties of the lattice $\ker_{\mathbb{Z}} \mathcal{A}_{\mathcal{C},\mathbf{d}}$, the polyhedral cone $\mathbb{R}_{\geq 0}\mathcal{A}_{\mathcal{C},\mathbf{d}} := \{\mathcal{A}_{\mathcal{C},\mathbf{d}}x : x \geq 0\}$ and the semigroup $\mathbb{N}\mathcal{A}_{\mathcal{C},\mathbf{d}} := \{\mathcal{A}_{\mathcal{C},\mathbf{d}}x : x \geq 0, x \text{ integral }\}$. For example, Markov bases of the lattice $\ker_{\mathbb{Z}} \mathcal{A}_{\mathcal{C},\mathbf{d}}$ can be used for goodness-of-fit tests of the log-linear model associated to $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ (see [5]). In the special case where $\mathbf{d} = \mathbf{2}$ the vector of all twos, we call $\mathcal{A}_{\mathcal{C},\mathbf{2}}$ a binary hierarchical model.

There are a number of "niceness" properties that an integral matrix $A \in \mathbb{Z}^{d\times n}$ could satisfy. Perhaps the strongest is unimodularity.

Definition 1.1. The following are equivalent definitions for an integral matrix A to be *unimodular*.

(1) The polyhedron $P_{A,b} = \{x \in \mathbb{R}^s : Ax = b, x \geq 0\}$ has all integral vertices for every $b \in \mathbb{N}A$.
(2) Suppose $\text{rank}A = d$. Then there exists a λ such that the determinants of all $d \times d$ submatrices of A are 0 or $\pm\lambda$.
(3) The entries in each circuit of A are either 0 or ± 1.
(4) The Graver basis of A consists of elements whose entries are either 0 or ± 1.

We elaborate on this definition in Section 2. A weaker property for the matrix A to satisfy is normality.

Definition 1.2. A matrix A is *normal* if the semigroup $\mathbb{N}A$ is saturated, that is $\mathbb{N}A = \mathbb{Z}A \cap \mathbb{R}_{\geq 0}A$.

In fact, every unimodular matrix A is normal. In previous work of Rauh and the second author [11], normality was identified as a key property of hierarchical models to

be able to apply the toric fiber product construction to calculate a Markov basis. If $A_{\mathcal{C},\mathbf{d}}$ is unimodular, it is also easy to solve the integer programs that arise in sequential importance sampling [4]. It is a major open problem to classify the normal hierarchical models. The special case where the underlying simplicial complex $\mathcal{C}$ is a graph, and where $\mathbf{d} = \mathbf{2}$ was handled in [16] where it was shown that a binary graph model is normal if and only if the graph is free of K_4 minors.

Our goal in the present paper is to classify the unimodular binary hierarchical models. In fact, this result is also an important first step in the classification of normal binary hierarchical models because a matrix $\Lambda(A)$ of Lawrence-type is normal if and only if unimodular (this follows from [10] Corollary 16 in light of the circuit definition of unimodularity). So this provides a characterization of the normal binary hierarchical models that have a big facet (see Definition 3.14 and remarks that follow).

Our main result is a complete structural characterization of the unimodular binary hierarchical models. On the one hand, we show that every unimodular binary hierarchical model can be built up from three basic families, together with three operations. As part of that proof we also give an excluded minor characterization of the unimodular binary hierarchical models. Here our notion of minor of a simplicial complex is one that is obtained by taking a sequence of vertex deletions or links of vertices. We caution the reader that our notion of simplicial complex minor does not directly generalize the notion of a graph minor, but it does generalize the notion of a matroid minor. Our excluded minor characterization involves an infinite number of minimally nonunimodular complexes.

Our characterization of unimodular binary hierarchical models can be seen in analogy with the classic characterizations of regular matroids/totally unimodular matrices, which have both excluded minor characterizations and constructive characterizations. While there is clearly a close connection between our problem and the study of totally unimodular matrices, our methods do not make use of Seymour's [13] nor Tutte's [17] classification of regular matroids. There are two reasons for this. The first is that the set of matroids underlying the design matrices of hierarchical models is not closed under the operations of direct sum, two sum, three sum, deletion, nor contraction. The second is that normality, unlike (total) unimodularity, is not a matroidal property and we are interested in extending our methods to the normal classification problem [2].

The outline of this paper is as follows. In the next section, we provide a detailed introduction to hierarchical models and the construction and interpretation of the matrix $\mathcal{A}_{\mathcal{C},\mathbf{d}}$. Section 3 details basic examples of unimodular complexes and constructions of new unimodular complexes from old ones. A main result here is Proposition 3.7 which shows that Alexander duality preserves unimodularity. Section 4 gives the list of minimal non-unimodular binary hierarchical models. There are one infinite family and six sporadic minimal non-unimodular complexes. The section also provides a number of structural results on simplicial complexes that avoid this list of simplicial complexes. Section 5 provides the combinatorial description of the 1-skeleton of a simplicial complex that supports a unimodular binary hierarchical model. That result is used as a stepping stone in a complex induction in Section 6 which gives the proof of the main theorem, Theorem

6.3. In Section 7 we explain how our characterization in the binary case can be used to make headway in the non-binary case.

2. Preliminaries on Hierarchical Models and Unimodularity

In this section we explain how to construct the matrix $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ associated to a hierarchical model. We then define some terms from Definition 1.1 of a unimodular matrix and prove the equivalence between the four parts.

Definition 2.1. Let $\mathbf{d} = (d_1, \ldots, d_n) \in \mathbb{Z}^n$ be such that $d_i \geq 2$ for each i. Then we define $\mathbb{R}^{\mathbf{d}}$ to be the vector space of all $d_1 \times \cdots \times d_n$-way tables. That is, each $\mathbf{u} \in \mathbb{R}^{\mathbf{d}}$ is a table in n dimensions where the ith dimension has d_i levels. We denote entries in a table $\mathbf{u}$ by $u_{i_1,\ldots,i_n}$ where $(i_1, \ldots, i_n) \in [d_1] \times \cdots \times [d_n]$. We will use the shorthand $\mathbf{i}$ to denote $i_1, \ldots, i_n$.

Definition 2.2. Let $\mathcal{C}$ be a simplicial complex on ground set $[n]$. A *facet* of $\mathcal{C}$ is an inclusion-maximal subset $F \subseteq [n]$ that is contained in $\mathcal{C}$. We let $\text{facet}(C)$ denote the collection of facets of $\mathcal{C}$.

Definition 2.3. Let $\mathbf{d} = (d_1, \ldots, d_n) \in \mathbb{Z}^n$ and $F = \{f_1, \ldots, f_k\} \subseteq [n]$. We define $\mathbf{d}_F$ to be the restriction of $\mathbf{d}$ to the indices in F, i.e. $\mathbf{d}_F = (d_{f_1}, \ldots, d_{f_k})$.

Definition 2.4. Let $\mathcal{C}$ be a simplicial complex on ground set $[n]$ and let $\mathbf{d} = (d_1, \ldots, d_n) \in \mathbb{Z}^n$ be such that $d_i \geq 2$ for each i. Then we have the linear map

$$\pi_{\mathcal{C},\mathbf{d}} : \mathbb{R}^{\mathbf{d}} \to \bigoplus_{F \in \text{facet}(C)} \mathbb{R}^{\mathbf{d}_F}$$

defined by

$$(u_{i_1,\ldots,i_n} : i_j \in [d_j]) \mapsto \bigoplus_{\substack{F \in \text{facet}(\mathcal{C}) \\ F = \{f_1,\ldots,f_k\}}} \left(\sum_{\substack{\mathbf{i} \in [d_1] \times \cdots \times [d_n]: \\ \mathbf{i}_F = \mathbf{j}}} u_{i_1,\ldots,i_n} : \mathbf{j} \in [d_{f_1}] \times \cdots \times [d_{f_k}] \right).$$

The matrix $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ denotes the matrix in the standard basis that represents the linear transformation $\pi_{\mathcal{C},\mathbf{d}}$.

We give an example of this construction.

Example 2.5. Let $\mathcal{C}$ be the simplicial complex on ground set $[3] = \{1, 2, 3\}$ with $\text{facet}(\mathcal{C}) = \{\{1\}, \{2, 3\}\}$. The linear transformation $\pi_{\mathcal{C},\mathbf{d}}$ maps a three-way tensor $u = (u_{ijk} : i \in [d_1], j \in [d_2], k \in [d_3]) \in \mathbb{R}^D$ to the direct sum of a one-way tensor and a two-way tensor:

$$\pi_{\mathcal{C},\mathbf{d}}(u) = (\sum_{j,k} u_{ijk} : i \in [d_1]) \oplus (\sum_{i} u_{ijk} : j \in [d_2], k \in [d_3]).$$

Taking $\mathbf{d} = \mathbf{2}$, the matrix $\mathcal{A}_{\mathcal{C},\mathbf{2}}$ could be represented as follows:

$$\left(\begin{array}{cccccccc} 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ \hline 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{array}\right),$$

the horizontal line dividing the image space into the parts corresponding to the two facets of $\mathcal{C}$.

It can be useful to be precise about how to write down the matrix $\mathcal{A}_{\mathcal{C},\mathbf{d}}$. The columns of the matrix $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ are indexed by elements of the set $\prod_{j\in[n]}[d_j]$ and the rows are indexed by pairs (F, e) where $F \in \text{facet}(\mathcal{C})$ and $e \in \prod_{j\in F}[d_j]$. The entry in position with row index (F, e) and column indexed by $i \in \prod_{j\in[n]}[d_j]$ is 1 if $e = (i_j : j \in F)$, otherwise it is equal to 0.

The binary case studied in this paper concerns the situation where all table dimensions are of size 2, that is $\mathbf{d} = \mathbf{2} = (2, 2, \ldots, 2)$; the vector of all twos. In this case, we will usually abbreviate notation as stated in the following definition.

Definition 2.6. Let $\mathcal{C}$ be a simplicial complex on $[n]$. We define $\mathcal{A}_{\mathcal{C}} := \mathcal{A}_{\mathcal{C},\mathbf{2}}$. We say that $\mathcal{C}$ is *unimodular* if $\mathcal{A}_{\mathcal{C}}$ is unimodular as a matrix.

We now elaborate on Definition 1.1. To start, we give two definitions.

Definition 2.7. Let A be an integer matrix. A nonzero element $u \in \ker_{\mathbb{Z}} A$ is called a *circuit* if its nonzero entries are relatively prime and there is no other nonzero element $v \in \ker_{\mathbb{Z}} A$ such that $\text{supp}(v) \subset \text{supp}(u)$. Here, $\text{supp}(v)$ denotes the set of positions of nonzero elements of v.

Definition 2.8. Let A be an integer matrix. A nonzero element $u \in \ker_{\mathbb{Z}} A$ is called *primitive* if there is no nonzero $v \in \ker_{\mathbb{Z}} A$, $v \neq u$ such that $u_i v_i \geq 0$ for all i, and $|v_i| \leq |u_i|$ for all i. The set of all primitive vectors in $\ker_{\mathbb{Z}} A$ is called the *Graver basis* of A.

Now we explain the equivalence between the four items in Definition 1.1. See [12], Theorem 19.2 for the equivalence of (1) and (2). We can see that condition (3) of Definition 1.1 is equivalent to condition (2) because the circuits can be computed via a determinantal formula using Cramer's rule [14, p. 35]. Given the equivalence of (2) and (3), (3) and (4) are equivalent by [14], Propositions 4.11 and 8.11.

Questions about unimodularity of a matrix A are invariant under certain changes to the matrix. The following proposition gives three that we make tacit use of throughout the rest of the paper.

Proposition 2.9. *Let A be a unimodular integer matrix. Then the following are also unimodular matrices.*

(1) *A matrix obtained by reordering the columns of A*

(2) *A matrix obtained by multiplying a column of A by -1*
(3) *Any matrix that has the same rowspace as A*
(4) *Any matrix formed by a subset of columns of A.*

Proof. Unimodularity of (1) and (3) are clear when considering (3) or (4) in Definition 1.1. Unimodularity of (2) is clear when considering (2) in Definition 1.1. If A' is a matrix obtained by extracting a set of columns of A then the circuits of A' are obtained by taking those circuits of A whose nonzero elements correspond to the columns of A we are extracting, and so (4) holds. □

3. Constructions of Unimodular Complexes

In this section we describe some operations on simplicial complexes that preserve unimodularity. In particular we show that unimodularity is preserved when passing to induced subcomplexes, to the Alexander dual of a simplicial complex, and to the link of a face of a simplicial complex. We can also build new unimodular complexes from old ones by adding cone vertices, ghost vertices, or taking a Lawrence lifting. We also construct the basic examples of unimodular complexes. These tools together go in to our constructive description of unimodular complexes.

Proposition 3.1. *All induced sub-complexes of a unimodular complex are unimodular.*

Proof. Let $\mathcal{C}'$ be the induced subcomplex of $\mathcal{C}$ obtained by restricting to some vertex set $F \subseteq [n]$. Let B be the matrix obtained by taking the columns of $\mathcal{A}_\mathcal{C}$ corresponding to the elements $i = (i_1, \ldots, i_n) \in \prod_{j\in[n]}[d_j]$ such that $i_j = 1$ for all $j \in [n] \setminus F$. Let B' be the matrix obtained from B by removing rows of all zeros. Then $B' = \mathcal{A}_{\mathcal{C}'}$. So $\mathcal{A}_{\mathcal{C}'}$ and B have the same Graver basis, which is a subset of the Graver basis of $\mathcal{A}_\mathcal{C}$. So unimodularity of $\mathcal{A}_\mathcal{C}$ implies unimodularity of $\mathcal{A}_{\mathcal{C}'}$. □

Next we will show that unimodularity is preserved under the Alexander duality operation.

Definition 3.2. Let $\mathcal{C}$ be a simplicial complex on $[n]$. The *Alexander dual* of $\mathcal{C}$, denoted $\mathcal{C}^*$ is defined as:

$$\mathcal{C}^* = \{S \subseteq [n] : [n] \setminus S \notin \mathcal{C}\}.$$

Note that if $\mathcal{C}$ has d faces, then $\mathcal{C}^*$ has $2^n - d$ faces and that the facets of $\mathcal{C}^*$ are the complements of the *minimal* non-faces of $\mathcal{C}$.

Since we are now restricting attention to the binary case, where $d_i = 2$ for each $i \in [n]$, entries of vectors in $\mathbb{R}^{\mathbf{d}}$ are indexed by elements of $\{1,2\}^n$ and entries of vectors in $\bigoplus_{f\in\text{facet}(\mathcal{C})} \mathbb{R}^{\mathbf{d}_F}$ are indexed by pairs $(S, \mathbf{j})$ where $S \in \text{facet}(\mathcal{C})$ and $\mathbf{j} \in \{1,2\}^S$. However, the statement and proof of Proposition 3.6 become cleaner if we replace $\{1,2\}^n$ and $\{1,2\}^S$ by $\{0,1\}^n$ and $\{0,1\}^S$, respectively. So we adapt this convention for the remainder of this section. We now introduce some notation that we use in Propositions 3.4 and 3.6.

Definition 3.3. Let $S \subseteq [n]$ and $\mathbf{j} \in \{0,1\}^S$ and $\mathbf{i} \in \{0,1\}^{[n]\setminus S}$. Then $e_{\mathbf{i},\mathbf{j}}$ denotes the element $v \in \mathbb{R}^{\{0,1\}^n}$ such that $v_\mathbf{k} = 1$ if $k_t = j_t$ when $t \in S$ and $k_t = i_t$ when $t \in [n] \setminus S$, and $v_\mathbf{k} = 0$ otherwise.

We are now ready to give an explicit description of $\mathcal{A}_{\mathcal{C}^*}$.

Proposition 3.4. *Let $\mathcal{C}$ be a simplicial complex on $[n]$. Let M be a matrix with the following set of columns:*

$$\left\{ \sum_{\mathbf{j}\in\{0,1\}^S} e_{\mathbf{i},\mathbf{j}} : S \text{ is a minimal non-face of } \mathcal{C}, \mathbf{i} \in \{0,1\}^{[n]\setminus S} \right\}.$$

Then $\mathcal{A}_{\mathcal{C}^*} = M^T$.

Proof. We index the columns of M by pairs $(S, \mathbf{i})$ where S is a minimal non-face of $\mathcal{C}$, and $\mathbf{i} \in \{0,1\}^{[n]\setminus S}$. The rows of M are indexed by elements of $\{0,1\}^n$. By construction, $M(\mathbf{k}, (S, \mathbf{i})) = 1$ if and only if $\mathbf{k}|_{[n]\setminus S} = \mathbf{i}$. Similarly, when F is a facet of $\mathcal{C}^*$, $\mathcal{A}_{\mathcal{C}^*}((F, \mathbf{i}), \mathbf{k}) = 1$ if and only $\mathbf{k}|_F = \mathbf{i}$. The result follows since S is a minimal non-face of $\mathcal{C}$ if and only if $[n] \setminus S$ is a facet of $\mathcal{C}^*$. □

In order to relate the unimodularity of $\mathcal{A}_{\mathcal{C}}$ and $\mathcal{A}_{\mathcal{C}^*}$, we need two propositions. The first is a standard result from the theory of matroid duality, though we could not find a precise reference.

Proposition 3.5. *Let $A \in \mathbb{K}^{r\times n}$ have rank r. Let $B \in \mathbb{K}^{(n-r)\times n}$ have rank $n - r$ such that $AB^T = 0$. Then there exists a non-zero scalar $\lambda \in \mathbb{K}^*$ such that for all $S \subseteq [n]$ with $\#S = r$,*

$$\det(A_S) = \pm\lambda \det(B_{[n]\setminus S}).$$

Proof. After multiplying by an element of $GL_r(\mathbb{K})$ and permuting columns, we can assume A has the form $A = (I_r\ M)$ where I_r is an $r \times r$ identity matrix. Then we can suppose B has the form $B = (-M^T\ I_{n-r})$. Then any $\det(A_S)$ is a minor of M with row indices $[r] \setminus S$ and column indices $S \setminus [r]$. The determinant $\det(B_{[n]\setminus S})$ is the same minor of M, up to a sign. □

Proposition 3.6. *Let $\mathcal{C}$ be a simplicial complex on $[n]$. Then the following set of vectors spans $\ker(\mathcal{A}_{\mathcal{C}})$:*

$$K = \left\{ \sum_{\mathbf{j}\in\{0,1\}^S} (-1)^{\|\mathbf{j}\|_1} e_{\mathbf{i},\mathbf{j}} : S \text{ is a minimal non-face of } \mathcal{C}, \mathbf{i} \in \{0,1\}^{[n]\setminus S} \right\}.$$

Furthermore, if we view the entries of K as the columns of a matrix M', then we can multiply some set of rows and columns of M' by -1 and get M, as defined in Proposition 3.4.

Proof. First, note that for any $x \in K$, $\mathcal{A}_{\mathcal{C}}x = 0$, so $K \subset \ker(\mathcal{A}_{\mathcal{C}})$. If we view the entries of K as the columns of a matrix M', we are done if we show that $\operatorname{rank}(M') = \dim(\ker(\mathcal{A}_{\mathcal{C}}))$. We proceed by proving the second statement of the lemma. The first statement will follow since Proposition 3.4 implies $\operatorname{rank}(M) = \operatorname{rank}(\mathcal{A}_{\mathcal{C}^*})$ and we know $\operatorname{rank}(\mathcal{A}_{\mathcal{C}^*}) = 2^n - \#\mathcal{C} = \dim(\ker(\mathcal{A}_{\mathcal{C}}))$ ([6, Thm 2.6]).

As before, we index the columns of M' by pairs $(S, \mathbf{i})$ where S is a minimal non-face of $\mathcal{C}$, and we index the rows of M' by the binary n-tuples. Let $M'_{(S,\mathbf{i})}$ be a column of M'.

For any $\mathbf{k} \in \{0,1\}^n$, the entry $M'_{(S,\mathbf{i}),\mathbf{k}}$ is nonzero if and only if $\mathbf{k}|_{[n]\setminus S} = \mathbf{i}$ - i.e. for each column, the $\mathbf{k}$ for each nonzero entry has fixed $\mathbf{k}|_{[n]\setminus S} = \mathbf{i}$. Now in this case, recall that $M'_{(S,\mathbf{i}),\mathbf{k}} = +1$ if $\|\mathbf{j}\|_1$ is even, and $M'_{(S,\mathbf{i}),\mathbf{k}} = -1$ if $\|\mathbf{j}\|_1$ is odd. Therefore, when $\|\mathbf{i}\|_1$ is even, $M'_{(S,\mathbf{i}),\mathbf{k}} = +1$ if $\|\mathbf{k}\|_1$ is even and $M'_{(S,\mathbf{i}),\mathbf{k}} = -1$ when $\|\mathbf{k}\|_1$ is odd. And when $\|\mathbf{i}\|_1$ is odd, $M'_{(S,\mathbf{i}),\mathbf{k}} = +1$ when $\|\mathbf{k}\|_1$ is odd and $M'_{(S,\mathbf{i}),\mathbf{k}} = -1$ when $\|\mathbf{k}\|_1$ is even. So for a fixed column $M'_{(S,\mathbf{i})}$, the sign of a nonzero entry $M'_{(S,\mathbf{i}),\mathbf{k}}$ depends only on the parity of $\mathbf{k}$. So if we multiply all the rows $M'_{\cdot,\mathbf{k}}$ such that $\|\mathbf{k}\|_1$ is odd by -1, the entries in each column will all have the same sign. Then we can multiply all the negative columns by -1, to arrive at the matrix from Proposition 3.4. □

Now we are ready to show that Alexander duality preserves unimodularity.

Proposition 3.7. *Let $\mathcal{C}$ be a simplicial complex on ground set $[n]$. Then $\mathcal{C}$ is unimodular if and only if $\mathcal{C}^*$ is unimodular.*

Proof. Consider the set K from Proposition 3.6 as a matrix of column vectors. Since K spans the kernel of $\mathcal{A}_\mathcal{C}$, Proposition 3.5 implies that K^T is unimodular if and only if $\mathcal{A}_\mathcal{C}$ is unimodular. Then we can multiply the appropriate rows and columns of K by -1 to get M as in Proposition 3.4. Since these operations preserve the absolute values of full rank determinants, K^T is unimodular if and only if M^T is. Proposition 3.4 says that $\mathcal{A}_{\mathcal{C}^*} = M^T$. □

Taking Alexander duals and induced complexes gives rise to another unimodularity preserving operation which we now define.

Definition 3.8. Let $S \in \mathcal{C}$ be a face of $\mathcal{C}$. Then the link of S in $\mathcal{C}$ is the new simplicial complex

$$\operatorname{link}_S(\mathcal{C}) = \{F \setminus S : F \in \mathcal{C} \text{ and } S \subseteq F\}.$$

When $S = \{v\}$, we simply write $\operatorname{link}_v(\mathcal{C}) := \operatorname{link}_{\{v\}}(\mathcal{C})$.

Note that we can obtain $\operatorname{link}_S(\mathcal{C})$ by repeatedly taking links with respect to vertices. That is if S is a face of $\mathcal{C}$ and $\#S \geq 2$ and $v \in S$, then

$$\operatorname{link}_S(\mathcal{C}) = \operatorname{link}_v(\operatorname{link}_{S\setminus\{v\}}(\mathcal{C})).$$

Proposition 3.9. *If $\mathcal{C}$ is a simplicial complex and S is a face of $\mathcal{C}$, then* $\operatorname{link}_S(\mathcal{C}) = (\mathcal{C}^* \setminus S)^*$.

Proof. By definition, we have:

$$(\mathcal{C}^* \setminus S)^* = \{R \subseteq [n] \setminus S : ([n] \setminus S) \setminus R \notin \mathcal{C}^* \setminus S\}.$$

Then we have the following chain of equivalences on some $R \subseteq [n] \setminus S$:

$$\begin{aligned} &([n] \setminus S) \setminus R \notin \mathcal{C}^* \setminus S \\ &\iff ([n] \setminus S) \setminus R \notin \mathcal{C}^* \\ &\iff [n] \setminus (([n] \setminus S) \setminus R) \in \mathcal{C} \end{aligned}$$

and since $[n] \setminus (([n] \setminus S) \setminus R) = R \cup S$, we have $R \cup S \in \mathcal{C}$. □

Corollary 3.10. *Let $\mathcal{C}$ be a unimodular simplicial complex on ground set $[n]$. Then for any face S of $\mathcal{C}$, $\text{link}_S(\mathcal{C})$ is unimodular.*

Proof. Proposition 3.9 implies that $\text{link}_S(\mathcal{C})$ can be obtained via Alexander duality and passing to an induced subcomplex. Unimodularity then follows from Propositions 3.1 and 3.7. □

Now we turn to operations for taking a complex that is unimodular and constructing larger unimodular complexes. If $\mathcal{C}$ has a vertex v that lies in each facet of $\mathcal{C}$, we say that v is a *cone vertex* of $\mathcal{C}$. The following proposition tells us that unimodularity is invariant under adding or removing a cone vertex.

Proposition 3.11. *Let $\mathcal{C}$ be a simplicial complex on $[n]$. We define $\mathcal{C}'$ on $[n+1]$ to be the simplicial complex with the following facets:*

$$\{F \cup \{n+1\} : F \text{ is a facet of } \mathcal{C}\}.$$

Then $\mathcal{A}_\mathcal{C}$ is unimodular if and only if $\mathcal{A}_{\mathcal{C}'}$ is.

Proof. We will index the columns of $\mathcal{A}_{\mathcal{C}'}$ by the binary $n+1$ tuples such that those with the $n+1$ coordinate equal to 1 come before those with $n+1$ coordinate equal to 2. Then we get the following block form:

$$\mathcal{A}_{\mathcal{C}'} = \left(\begin{array}{c|c} \mathcal{A}_\mathcal{C} & \mathbf{0} \\ \hline \mathbf{0} & \mathcal{A}_\mathcal{C} \end{array}\right).$$

The Graver basis of $\mathcal{A}_{\mathcal{C}'}$ is therefore $\{(u,0),(0,u) : u \in Gr_A\}$. Hence the Graver basis of $\mathcal{A}_{\mathcal{C}'}$ consists of $0, \pm 1$ elements if and only if the Graver basis of $\mathcal{A}_\mathcal{C}$ consists of $0, \pm 1$ elements. □

By induction, adding or removing multiple cone vertices from a simplicial complex does not affect unimodularity. We introduce the following notation to denote this.

Definition 3.12. Let $\mathcal{C}$ be a simplicial complex on vertex set $[n]$. Then we define $\text{cone}^p(\mathcal{C})$ to be the simplicial complex on $[n] \cup \{u_1, \ldots, u_p\}$ with the following facets

$$\{F \cup \{u_1, \ldots, u_p\} : F \text{ is a facet of } \mathcal{C}\}.$$

When $p = 0$, we define $\text{cone}^p(\mathcal{C}) = \mathcal{C}$.

Definition 3.13. For any matrix $A \in \mathbb{R}^{s \times t}$, we define the *Lawrence lifting* of A to be the matrix

$$\Lambda(A) \quad = \quad \begin{pmatrix} A & \mathbf{0} \\ \mathbf{0} & A \\ \mathbf{1} & \mathbf{1} \end{pmatrix} \quad \in \quad \mathbb{R}^{(2s+t) \times 2t}$$

where $\mathbf{0}$ denotes the $s \times t$ matrix of all zeroes and $\mathbf{1}$ denotes a $t \times t$ identity matrix.

By Theorem 7.1 in [14], $\Lambda(A)$ is unimodular if and only if A is unimodular. This gives rise to another unimodularity-preserving operation on simplicial complexes.

Definition 3.14. Let $\mathcal{C}$ be a simplicial complex on $[n]$. We define the *Lawrence lifting* of $\mathcal{C}$ to be the simplicial complex $\Lambda(\mathcal{C})$ on $[n+1]$ that has the following set of facets:

$$\{[n]\} \cup \{F \cup \{n+1\} : F \text{ is a facet of } \mathcal{C}\}.$$

In this case, we refer to the facet $[n]$ as a *big facet*.

If a complex $\mathcal{C}$ on $[n+1]$ has big facet $[n]$, then $\mathcal{C} = \Lambda(\text{link}_{n+1}(\mathcal{C}))$. In particular, $\mathcal{C} = \Lambda(\mathcal{D})$ for some complex $\mathcal{D}$ if and only if $\mathcal{C}$ has a big facet.

Proposition 3.15. *The simplicial complex $\mathcal{C}$ is unimodular if and only if $\Lambda(\mathcal{C})$ is unimodular.*

Proof. Note that by construction $\Lambda(\mathcal{A}_{\mathcal{C}}) = \mathcal{A}_{\Lambda(\mathcal{C})}$. Hence Theorem 7.1 in [14] implies the proposition. □

If a simplicial complex $\mathcal{C}$ can be expressed as $\Lambda(\mathcal{C}')$, then we say that $\mathcal{C}$ is of Lawrence type. A simplicial complex on n vertices is of Lawrence type if and only if it has a facet containing $n-1$ vertices. We will refer to any facet of $\mathcal{C}$ that has $n-1$ elements as a *big facet*.

We now give our final unimodularity preserving operation.

Definition 3.16. Let $\mathcal{C}$ be a simplicial complex on ground set $[n]$. Let $G^p(\mathcal{C})$ denote the same simplicial complex but on ground set $[n+p]$. Note that the vertices $n+1, \ldots, n+p$ are not contained in any face of $G^p(\mathcal{C})$. In this case we say that $n+1, \ldots, n+p$ are *ghost vertices*. When $p = 1$ we drop the superscript; i.e. we just write $G(\mathcal{C})$.

Proposition 3.17. *A simplicial complex $\mathcal{C}$ is unimodular if and only if $G^p(\mathcal{C})$ is unimodular.*

Proof. This is true because

$$\mathcal{A}_{G^p(\mathcal{C})} = \begin{pmatrix} \mathcal{A}_{\mathcal{C}} & \ldots & \mathcal{A}_{\mathcal{C}} \end{pmatrix}.$$ □

Note that adding a ghost vertex to a complex is Alexander dual to taking the Lawrence lifting of a simplicial complex, that is $G(\mathcal{C})^* = \Lambda(\mathcal{C}^*)$.

We now state a useful fact about the interaction of these operations.

Proposition 3.18. *The operation* $\text{cone}^p(\cdot)$ *commutes with the operations of taking links, duals, induced sub-complexes and adding ghost vertices.*

For any vertex v of $\mathcal{C}$, we let $\mathcal{C} \setminus v$ denote the induced subcomplex on $\mathcal{C} \setminus \{v\}$. Then for vertices $v \neq u$ of $\mathcal{C}$, the operations $\cdot \setminus v$ and $\text{link}_u(\cdot)$ commute. So if $\mathcal{D}$ can be obtained from $\mathcal{C}$ by applying a series of deletions, and taking links, then we can write

$$\mathcal{D} = \text{link}_R(\mathcal{C} \setminus S)$$

where $F \in \mathcal{C}$ and S, R are a subset of vertices of $\mathcal{C}$ such that $S \cap R = \emptyset$. This gives rise to the following definition:

Definition 3.19. Let $\mathcal{C}$ be a simplicial complex. Then $\mathcal{D}$ is a *minor* of $\mathcal{C}$ if $\mathcal{D} = \text{link}_R(\mathcal{C} \setminus S)$ where and S, R are subsets of vertices of $\mathcal{C}$ such that $S \cap R = \emptyset$, and R is a face of $\mathcal{C}$.

We warn the reader that although it is natural to view simplicial complexes as a generalization of graphs, our definition of simplicial complex minor does *not* generalize the usual notion of graph minor. Alternatively, a simplicial complex can be seen as a generalization of a matroid, and our definition of minor is a generalization of the usual notion of a matroid minor [8]. This definition of simplicial complex minor is useful for our purposes because of Proposition 3.20.

Proposition 3.20. *If $\mathcal{C}$ is a unimodular simplicial complex, then every minor of $\mathcal{C}$ is unimodular.*

Proof. This follows immediately from Propositions 3.1 and 3.10. □

The fundamental example of a unimodular simplicial complex is the disjoint union of two simplices. Here a simplex Δ_n is the simplicial complex on an $n+1$ element set with a single facet consisting of all the elements. We denote the irrelevant complex $\{\emptyset\}$ by Δ_{-1} and the void complex $\{\}$ by Δ_{-2}.

Proposition 3.21. *Let $\mathcal{C} = \Delta_m \sqcup \Delta_n$ be the disjoint union of two simplices, for $m, n \geq 0$. Then $\mathcal{C}$ is unimodular.*

Proof. The matrix $\mathcal{A}_\mathcal{C}$ in this case is the vertex-edge incidence matrix of a complete bipartite graph with 2^{m+1} and 2^{n+1} vertices in the two parts of the partition. Such vertex-edge incidence matrices are examples of network matrices and are hence totally unimodular [12, Ch. 19]. □

A second family of fundamental examples comes from taking the duals of the disjoint union of two simplices. We use $D_{m,n}$ to denote the dual of a disjoint union of an m-simplex and an n-simplex, i.e.

$$D_{m,n} := (\Delta_m \sqcup \Delta_n)^*.$$

We close this section by giving a workable description of $D_{m,n}$. We can divide the vertices of $D_{m,n}$ into disjoint sets M, N such that M contains the vertices of the Δ_m in $D^*_{m,n}$ and N contains the vertices of the Δ_n in $D^*_{m,n}$. Then, the facets of $D_{m,n}$ are precisely the subsets of $M \sqcup N$ that leave out exactly one element of M and one element of N. Notice that the complexes induced on M, N are $\partial\Delta_m$ and $\partial\Delta_n$ respectively. Also, note that for any $v \in M$, $\text{link}_v(D_{m,n}) = D_{m-1,n}$ and for any $v \in N$, $\text{link}_v(D_{m,n}) = D_{m,n-1}$.

4. β-avoiding Simplicial Complexes

Part of our main result is a forbidden minor classification of unimodular simplicial complexes. In this section, we identify these forbidden minors and prove various properties about the complexes that avoid them.

Proposition 4.1. *The following simplicial complexes are minimal nonunimodular simplicial complexes:*

1. *P_4, the path on 4 vertices*
2. *O_6, the boundary of the octahedron or its dual O_6^**
3. *J_1, the complex on $\{1,2,3,4,5\}$ with facets $12, 15, 234, 345$ or its dual J_1^*, with facets $134, 235, 245$.*

(4) J_2, *complex on* $\{1,2,3,4,5\}$ *with facets* $12, 235, 34, 145$
(5) *For* $n \geq 1$, $\partial\Delta_n \sqcup \{v\}$, *the disjoint union of the boundary of an* n*-simplex and a single vertex.*




FIGURE 1. The complexes P_4, J_1, J_1^*, and J_2.

Note that P_4, J_2 and $\partial\Delta_n \sqcup \{v\}$ are isomorphic to their own duals.

Proof. We can check that the complexes $P_4, O_6, O_6^*, J_1, J_1^*$, and J_2 are not unimodular by using the software 4ti2 [1] to compute the Graver basis and looking for entries that are not $0, \pm 1$ In the case of O_6 and O_6*, these are too large to compute the entire Graver basis. However, selecting sufficiently large random subsets of the columns produced Graver basis elements of the desired form. Computations are available on our website [3]. For the infinite family $\partial\Delta_n \sqcup \{v\}$ where $n \geq 1$, examples of a non-squarefree Graver basis element appear in [15]. These results show that these examples are not unimodular. To see that they are minimal, note that every subcomplex obtained by deleting a single vertex or taking the link at a vertex produced a unimodular complex in all cases. □

Definition 4.2. A simplicial complex $\mathcal{C}$ is *β-avoiding* if it does not contain any of the complexes from Proposition 4.1 as minors.

Since none of the complexes from Proposition 4.1 is unimodular, Proposition 3.20 implies that if $\mathcal{C}$ is unimodular, then $\mathcal{C}$ is β-avoiding. The converse of this is our forbidden minor classification of unimodular complexes which we prove in Theorem 6.3.

We now give some technical results about β-avoiding complexes. Before beginning, we remind the reader that $\partial\Delta_1 \sqcup \{v\}$ is an independent set on 3 vertices, and so no β-avoiding complex can have an independent set of size 3.

Proposition 4.3. *Let* $\mathcal{C}$ *be a* β*-avoiding simplicial complex. Then* $\mathcal{C}^*$ *is also* β*-avoiding.*

Proof. The list of prohibited minors of β-avoiding complexes is closed under taking duals. The proposition follows when we note that if $\mathcal{D}$ is a minor of $\mathcal{C}$, then $\mathcal{D}^*$ is a minor of $\mathcal{C}^*$. This is true because if $\mathcal{D}^* = \text{link}_S(\mathcal{C}^* \setminus R)$, then two applications of Proposition 3.9 give

$$\begin{aligned} \mathcal{D}^* &= ((\mathcal{C}^* \setminus R)^* \setminus S)^* \\ &= (\text{link}_R(\mathcal{C}) \setminus S)^* \end{aligned}$$

and therefore $\mathcal{D} = \text{link}_R(\mathcal{C}) \setminus S$. So $\mathcal{D}$ is a minor of $\mathcal{C}$. □

Proposition 4.4. *Let* $\mathcal{C}$ *be a* β*-avoiding simplicial complex that has* C_4 *induced. Then the complex induced on the non-ghost vertices of* $\mathcal{C}$ *is* $\text{cone}^p(C_4)$ *for some* p*.*

Proof. Assume $\mathcal{C}$ has no ghost vertices. Let $u_1, \ldots, u_4$ be the vertices that induce C_4 in $\mathcal{C}$. If these are the only vertices, we are done, so let v be another vertex in $\mathcal{C}$. Let $\mathcal{D}$ denote the complex induced on $v, u_1, \ldots, u_4$. Any minor of $\mathcal{D}$ is a minor of $\mathcal{C}$, so $\mathcal{D}$ must also be β-avoiding. Vertex v cannot be disconnected from $u_1, \ldots, u_4$ since otherwise $\mathcal{D}$ has an independent set of size 3. Furthermore, v must connect to $u_1, \ldots, u_4$ with a triangle. Otherwise if v connected to u_i with an edge, then u_i has edge degree 3 and so $\text{link}_v(\mathcal{D})$ is an independent set on 3 vertices. There are 4 possible triangles that v can join with $u_1, \ldots, u_4$ in $\mathcal{D}$ so that $u_1, \ldots, u_4$ induce C_4. If v is only in one such triangle, wlog $\{v, u_1, u_2\}$, then $\{v, u_1, u_3, u_4\}$ is an induced P_4. The two non-isomorphic complexes on 5 vertices that have two triangles and an induced C_4 are J_1 and J_2. If $\mathcal{D}$ has 3 of the possible triangles, then $\text{link}_v(\mathcal{D})$ is P_4. So $\mathcal{D}$ must have all four - i.e. it must be $\text{cone}^1(C_4)$.

Any vertices $v, v' \notin \{u_1, u_2, u_3, u_4\}$ must connect to each other. Otherwise, the induced complex on $v, v', u_1, \ldots, u_4$ is O_6 which is not unimodular. Now we denote the vertices of $\mathcal{C}$ that are not any of the u_is as $v_1, \ldots, v_k$. Since v_k is a cone point over the C_4 on $u_1, \ldots, u_4$ and since v_k connects to all $v_j, j < k$, $\text{link}_{v_k}(\mathcal{C})$ is a β-avoiding complex on $v_1, \ldots, v_{k-1}, u_1, \ldots, u_4$ that has C_4 induced on $u_1, \ldots, u_4$. So, by induction on the number of vertices, $\text{link}_{v_k}(\mathcal{C})$ is an iterated cone over C_4, $\text{cone}^{k-1}(C_4)$. But also the induced complex on $v_1, \ldots, v_{k-1}, u_1, \ldots, u_4$ is $\text{cone}^{k-1}(C_4)$ by induction. This implies that $\mathcal{C} = \text{cone}^k(C_4)$. □

Proposition 4.5. *Let $\mathcal{C}$ be a β-avoiding complex that has $D_{m,n}$ induced for some $m, n \geq 1$. Then the complex induced on the non-ghost vertices of $\mathcal{C}$ is $\text{cone}^p(D_{m,n})$ for some p.*

Proof. Assume $\mathcal{C}$ has no ghost vertices. Just as in the remarks at the end of section 3, we divide the vertices of $D_{m,n}$ into disjoint sets M, N such that M contains the vertices of the Δ_m in $D^*_{m,n}$ and N contains the vertices of the Δ_n in $D^*_{m,n}$. Now, we proceed by induction on m and n. Note that $D_{1,1} = C_4$, so the base case is handled by Proposition 4.4. Assume that, without loss of generality, $m \geq 2$.

Let $v \in \mathcal{C} \setminus D_{m,n}$. The vertex v must connect to some $u \in M$ to avoid inducing $\partial\Delta_m \sqcup \{v\}$. Then, $\text{link}_u(\mathcal{C})$ contains v and $D_{m-1,n}$, and so by induction, the complex induced on v and the vertices of $D_{m-1,n}$ in $\text{link}_u(\mathcal{C})$ is a cone over $D_{m-1,n}$. This means that v is in every facet that contains u. Since $m \geq 2$, u is connected to every other vertex in $D_{m,n}$ and thus v is attached to every vertex in $D_{m,n}$. Now we apply the same argument to each of the vertices in the set M to see that v is in every facet that contains any vertex in set M. Every facet of $D_{m,n}$ contains some element of M. So this implies that the induced complex on v and $D_{m,n}$ must be the cone over $D_{m,n}$.

Now assume v, v' are both vertices in $\mathcal{C} \setminus D_{m,n}$. If they were not connected by an edge, then the induced complex on v, v' and $D_{m,n}$ contains a minor which is isomorphic to O_6. This means that v, v' are connected by an edge. Taking the link $\text{link}_v(\mathcal{C})$ produces a smaller complex with an induced $D_{m,n}$ hence it must be a cone by induction on the number of vertices not in the $D_{m,n}$. This reduces us the case where $\text{link}_v(\mathcal{C})$ is $\text{cone}^{p-1}(D_{m,n})$ and $\mathcal{C} \setminus v$ is $\text{cone}^{p-1}(D_{m,n})$ which implies that $\mathcal{C}$ is $\text{cone}^p(D_{m,n})$. □

Proposition 4.6. *Let $\mathcal{C}$ be a β-avoiding simplicial complex on vertices $u_1, \ldots, u_{k+1}, v$ such that the complex induced on $\{u_1, \ldots, u_{k+1}\}$ is $\partial\Delta_k$. Then v must be in a k-simplex with some subset of the u_i's.*

Proof. We proceed by induction on k. For the base case $k = 1$, note that $\partial\Delta_1$ is two isolated points. In this case, v must connect to u_1 or u_2 as a 1-simplex (an edge) to avoid inducing $\partial\Delta_1 \sqcup \{v\}$.

Now assume $k > 1$. The vertex v must attach to some u_i to avoid inducing $\partial\Delta_k \sqcup \{v\}$. Then, $\text{link}(u_i)$ has v and $\partial\Delta_{k-1}$, so by induction, v must form a $k-1$ simplex with some collection $u_1, \ldots, \hat{u_i}, \ldots, \hat{u_j}, \ldots, u_{k+1}$. So in $\mathcal{C}$, v must be in a k simplex with $u_1, \ldots, \hat{u_j}, \ldots, u_{k+1}$. □

Proposition 4.7. *Let $\mathcal{C}$ be a β-avoiding simplicial complex on $m+n+2$ vertices that does not have a facet of dimension $m+n$. Assume $D_{m,n} \subseteq \mathcal{C}$. Then $\mathcal{C} = D_{m,n}$.*

Proof. Note that $\mathcal{C}^* \subseteq \Delta_m \sqcup \Delta_n$ because $D_{m,n} \subseteq \mathcal{C}$. Since $\mathcal{C}$ does not have a facet of dimension $m+n$, $\mathcal{C}^*$ has no ghost vertices. Since $\mathcal{C}^*$ has no ghost vertices, if $\mathcal{C}^* \subsetneq \Delta_m \sqcup \Delta_n$, then $\mathcal{C}^*$ would have an induced $\{v\} \sqcup \partial\Delta_k$ for some $k \geq 1$. In this case $\mathcal{C}^*$ would not be β-avoiding and so by Proposition 4.3, neither would $\mathcal{C}$. □

5. The 1-Skeleton of a β-avoiding Complex

In this section we prove Lemma 5.3 which gives a complete characterization of the 1-skeleton of a β-avoiding simplicial complex. This is a crucial technical lemma in the proof of Theorem 6.3. We start with a technical proposition about graphs.

Proposition 5.1. *Let H be a connected graph that avoids K_3 and P_4 as induced subgraphs. Then H is a complete bipartite graph.*

Proof. Let $u \in V(H)$, let $N(u)$ denote the neighbors of u, and let $M(u)$ denote the non-neighbors of u (this set includes u). The bipartition of the vertices of H will be $M(u)$ and $N(u)$. Let $v \in M(u) \backslash \{u\}$. Since H is connected, there exists a path $u = u_1, u_2, \ldots, u_k = v$. Assume k is minimal. Since $v \neq u$, $k > 1$. We cannot have $k = 2$ since u, v are non-neighbors. We cannot have $k = 4$, since in order to avoid an induced P_4, we would need an edge (u_i, u_{i+2}) inducing a K_3, or an edge (u, v) contradicting that u, v are non-neighbors. If $k \geq 5$, there must exist an edge (u_1, u_4) to avoid an induced P_4, contradicting minimality of k. So we have $k = 3$. So for any $v \in M(u) \setminus \{u\}$, there exists a path u, a, v.

Now we show that H is bipartite with bipartition $M(u)$ and $N(u)$. It is clear that $N(u)$ is an independent set of vertices, for if $v, w \in N(u)$ had an edge between them, there would be a K_3 on u, v, w. Now we show that $M(u)$ is an independent set of vertices. Assume $w, v \in M(u)$. If either w, v is u, there is no edge between them by definition of $M(u)$, so assume $w, v \neq u$. Then by the above, we have paths u, a, v and u, b, w. So an edge (v, w) would induce the 5-cycle u, a, v, w, b, u. This 5-cycle must have a chord to avoid inducing P_4, but any chord in a 5-cycle induces a K_3.

Now we show that H is complete bipartite. Let $x \in M(u)$ and $y \in N(u)$. Since H is connected, there is a path $x = u_1, ..., u_k = y$. We may without loss of generality assume $k \leq 3$ since otherwise we could shorten the path using the edge (u_1, u_4) required to avoid

a P_4. The sets $M(u)$ and $N(u)$ are disjoint, so $k \neq 1$. If $k = 3$, then there is an induced P_4 x, u_2, y, u. So $k = 2$, and so (x, y) is an edge. □

For a graph G we define the complement graph G^c on the same set of vertices such that (u, v) is an edge of G^c if and only if (u, v) is not an edge in G. Now we can use Proposition 5.1 to give a strong restriction on the structure of G^c whenever G is the 1-skeleton of a β-avoiding complex.

Proposition 5.2. *If G is the 1-skeleton of a β-avoiding simplicial complex $\mathcal{C}$, then each connected component of G^c is complete bipartite.*

Proof. Note that $\{v\} \sqcup \partial\Delta_1$ consists of three disconnected vertices, its complement graph is K_3. The path P_4 is its own complement. Since $\mathcal{C}$ is β-avoiding, G avoids P_4 and an independent set of size three as induced subgraphs. So each connected component of G^c avoids P_4 and K_3 as induced subgraphs. Proposition 5.1 therefore implies that each connected component of G^c is complete bipartite. □

Now we are ready to characterize the 1-skeleton of a β-avoiding simplicial complex.

Lemma 5.3. *Let $\mathcal{C}$ be a β-avoiding simplicial complex and let G denote its 1-skeleton. Then G is one of the following*

(a) K_N
(b) *Two complete graphs glued along a (possibly empty) common clique*
(c) *The iterated cone over a 4-cycle.*

Proof. By Proposition 5.2, we know that each connected component of G^c is complete bipartite. Let H denote the induced subgraph of G^c that removes all isolated vertices. If H is empty, then G^c is an independent set of vertices and therefore $G = K_N$. So assume H is nonempty. We claim that if H is neither $K_{m,n}$ nor $K_2 \sqcup K_2$ for $m, n \geq 1$, then H induces either $K_2 \sqcup K_2 \sqcup K_2$ or $P_3 \sqcup K_2$. To prove the claim, first assume that H avoids $K_2 \sqcup K_2 \sqcup K_2$. Since H is not $K_{m,n}$ and has no isolated vertices, this implies that H has exactly two components. Since H is not $K_2 \sqcup K_2$, some connected component of H has at least three vertices. So we have a P_3 induced by this component, and K_2 induced by the other component and so the claim is proven.

The complement graphs of $K_2 \sqcup K_2 \sqcup K_2$ or $P_3 \sqcup K_2$ are shown in Figure 2. Both graphs have C_4 induced, but neither is the 1-skeleton for an iterated cone over C_4 since neither graph has a suspension vertex. Proposition 4.4 therefore implies that neither is the 1-skeleton of a β-avoiding simplicial complex and so G may not induce either. So G^c may not induce $K_2 \sqcup K_2 \sqcup K_2$ nor $P_3 \sqcup K_2$ and therefore neither may H. The claim then implies that H must be either $K_{m,n}$ or $K_2 \sqcup K_2$. Assume G^c has p isolated vertices. If $H = K_{m,n}$ then G is a K_{m+p} and a K_{n+p} glued along a common K_p. If $H = K_2 \sqcup K_2$ then G is an iterated cone over a 4-cycle. □

As we see in the following proposition, the 1-skeleton of $\mathcal{C}$ completely determines $\mathcal{C}$ when it is obtained by gluing two complete graphs along an empty clique.

Proposition 5.4. *Let $\mathcal{C}$ be a β-avoiding simplicial complex such that the 1-skeleton of $\mathcal{C}$ is the disjoint union of cliques K_m and K_n. Then $\mathcal{C} = \Delta_m \sqcup \Delta_n$.*




FIGURE 2. The complement graphs of $P_3 \sqcup K_2$ and $K_2 \sqcup K_2 \sqcup K_2$

Proof. Let $v_1, \ldots, v_k$ be vertices of the K_m. If $\{v_1, \ldots, v_k\}$ is a minimal non-face of $\mathcal{C}$, then if we let u be a vertex in the K_n, then the complex indued on $v_1, \ldots, v_k, u$ is $\partial\Delta_k \sqcup \{u\}$. □

6. The Main Theorem

The goal of this section is to give a proof of Theorem 6.3 which gives a complete characterization of the unimodular binary hierarchical models.

We begin this section by defining a *nuclear* complex. A nuclear complex is a complex that can be obtained from a disjoint union of two simplices by adding cone vertices, adding ghost vertices, taking Lawrence liftings and taking Alexander duals. Since $\Delta_m \sqcup \Delta_n$ is unimodular and these operations all preserve unimodularity, nuclear complexes are unimodular. Part of our main result is the converse - unimodular complexes are nuclear.

Definition 6.1. A simplicial complex $\mathcal{C}$ is *nuclear* if one of the following is true

(1) $\mathcal{C} = \Lambda(\mathcal{D})$ where $\mathcal{D}$ is nuclear
(2) $\mathcal{C} = G(\mathcal{D})$ where $\mathcal{D}$ is nuclear
(3) $\mathcal{C} = \text{cone}^p(\Delta_m \sqcup \Delta_n)$ for $p, m, n \geq 0$
(4) $\mathcal{C} = \text{cone}^p(D_{m,n})$ for $p \geq 0$ and $m, n \geq 1$
(5) $\mathcal{C} = \Delta_k$ for $k \geq -2$.

Every nuclear complex $\mathcal{C}$ can be constructed by applying the operations $\text{cone}^p(\cdot)$, $G(\cdot)$, and $\Lambda(\cdot)$ to a complex $\mathcal{D}$ where $\mathcal{D}$ is of the form $\Delta_m \sqcup \Delta_n$, $D_{m,n}$, or Δ_k. We refer to $\mathcal{D}$ as the *nucleus* of $\mathcal{C}$.

Note that $D_{m,0}$ has a ghost vertex for all m. This is why we have $m, n \geq 1$ in (4). We note that the collection of nuclear complexes is closed under Alexander duality.

Proposition 6.2. *If $\mathcal{C}$ is nuclear then so is $\mathcal{C}^*$.*

Proof. Possibilities (1) and (2) are dual to each other, as are (3) and (4). On k vertices, Δ_k and Δ_{-2} are dual to each other. □

We now state our main result.

Theorem 6.3. *Let $\mathcal{C}$ be a simplicial complex. Then the following are equivalent*

(1) *$\mathcal{C}$ is unimodular*
(2) *$\mathcal{C}$ is β-avoiding*
(3) *$\mathcal{C}$ is nuclear.*

We defer the proof of Theorem 6.3 until the end of the section, but we give a roadmap here. It is immediate from results in previous sections that nuclear complexes are unimodular and that unimodular complexes are β-avoiding.

We show that any β-avoiding complex $\mathcal{C}$ is nuclear by choosing a particular vertex v of $\mathcal{C}$ and using induction on the number of vertices to conclude that $\text{link}_v(\mathcal{C})$ is nuclear. From here, we have five cases to consider - one for each of the ways that $\text{link}_v(\mathcal{C})$ can be nuclear. In four of these cases, it is relatively easy to show that $\mathcal{C}$ is nuclear. The difficulty lies in the case where $\text{link}_v(\mathcal{C}) = D_{m,n}$, which is handled in Lemma 6.8. The proof of Lemma 6.8 is split into two main parts and the second part is further split into six cases. Proposition 6.4 helps with the first part. Propositions 6.5, 6.6, and 6.7 handle the hard cases in the second part.

Proposition 6.4. *Let $\mathcal{C}$ be a β-avoiding simplicial complex. Assume that all proper minors of $\mathcal{C}$ are nuclear. Assume $\mathcal{C}$ has a vertex v such that $\mathcal{C} \setminus v = \text{cone}^p(\Delta_m \sqcup \Delta_n)$ with $m, n \geq 1$ and $p \geq 1$. Assume $\mathcal{C}$ is C_4-free. Then $\mathcal{C} = \text{cone}^{p+1}(\Delta_m \sqcup \Delta_n)$, $\mathcal{C} = \text{cone}^p(\Delta_{m+1} \sqcup \Delta_n)$, or $\mathcal{C} = \text{cone}^p(\Delta_m \sqcup \Delta_{n+1})$.*

Proof. We induct on p. For the base case, take $p = 1$. If we let u denote the cone vertex in $\mathcal{C} \setminus v$ then v must connect to one of the simplices in the $\Delta_m \sqcup \Delta_n \subset \mathcal{C} \setminus u$. Otherwise we have an independent set with three vertices induced in $\mathcal{C} \setminus u$ and this complex is not nuclear. If v connects to only one such simplex, then v must also connect to u to avoid inducing P_4 which is not nuclear. If v connects to both such simplices, and v does not connect to u, then $\mathcal{C}$ has C_4 induced which contradicts our hypothesis. So assume v connects to u and to at least one simplex. Since u is a cone vertex in $\mathcal{C} \setminus v$, $\text{link}_u(\mathcal{C})$ has no ghost vertices and by hypothesis, it is nuclear. Furthermore, $\text{link}_u(\mathcal{C} \setminus v) = \Delta_m \sqcup \Delta_n$ and so $\text{link}_u(\mathcal{C})$ is either $\Delta_{m+1} \sqcup \Delta_n$ or $\text{cone}^1(\Delta_m \sqcup \Delta_n)$ (note that these are the only nuclear complexes that become disconnected upon removing a vertex). In the first case, $\mathcal{C} = \text{cone}^1(\Delta_{m+1} \sqcup \Delta_n)$ and in the second case $\mathcal{C} = \text{cone}^2(\Delta_m \sqcup \Delta_n)$.

Now assume $p \geq 2$. Let $u_1, \ldots, u_p$ denote the cone vertices of $\mathcal{C} \setminus v$. Let M, N denote the vertex sets of the Δ_m, Δ_n respectively. Then v must connect to at least one of M and N to avoid inducing an independent set on three vertices. Furthermore, v must connect to each u_i - if v only connects to one of M or N, then this is required to avoid P_4, and if v connects to both then we need this to ensure that $\mathcal{C}$ is C_4-free. Furthermore, the set $\{u_1, \ldots, u_p, v\}$ is a facet of $\mathcal{C}$. This is clear when $p = 1$ since otherwise v doesn't connect to u_1. Induction on p gives that $\{u_1, \ldots, u_{p-1}, v\}$ is a facet of $\text{link}_{u_p}(\mathcal{C})$ and therefore $\{u_1, \ldots, u_p, v\}$ is a facet of $\mathcal{C}$.

Now, we can see that $\text{link}_{u_p}(\mathcal{C})$ has no ghost vertices and is C_4-free. Furthermore, $\text{link}_{u_p}(\mathcal{C}) \setminus v$ and $\mathcal{C} \setminus \{u_p, v\}$ are both equal to $\text{cone}^{p-1}(\Delta_m \sqcup \Delta_n)$. So by induction on p, each of $\text{link}_{u_p}(\mathcal{C})$ and $(\mathcal{C} \setminus u_p)$ can either be $\text{cone}^p(\Delta_m \sqcup \Delta_n)$ or $\text{cone}^{p-1}(\Delta_{m+1} \sqcup \Delta_n)$. If $\text{link}_{u_p}(\mathcal{C}) = \mathcal{C} \setminus u_p$, then u_p is a cone vertex in $\mathcal{C}$. So if both equal $\text{cone}^p(\Delta_m \sqcup \Delta_n)$ then $\mathcal{C} = \text{cone}^{p+1}(\Delta_m \sqcup \Delta_n)$. If both equal $\text{cone}^{p-1}(\Delta_{m+1} \sqcup \Delta_n)$ then $\mathcal{C} = \text{cone}^p(\Delta_{m+1} \sqcup \Delta_n)$.

Since $\text{link}_{u_p}(\mathcal{C}) \subseteq \mathcal{C} \setminus u_p$, it is impossible to have $\text{link}_{u_p}(\mathcal{C}) = \text{cone}^p(\Delta_m \sqcup \Delta_n)$ and $\mathcal{C} \setminus u_p = \text{cone}^{p-1}(\Delta_{m+1} \sqcup \Delta_n)$. It is also impossible to have $\mathcal{C} \setminus u_p = \text{cone}^p(\Delta_m \sqcup \Delta_n)$ and $\text{link}_{u_p}(\mathcal{C}) = \text{cone}^{p-1}(\Delta_{m+1} \sqcup \Delta_n)$. For if this were the case, then v would be a cone vertex in $\mathcal{C} \setminus u_p$ and it would be part of the Δ_{m+1} in $\text{link}_{u_p}(\mathcal{C})$. Since $n \geq 1$,

$\#N \geq 2$, so given $b_1, b_2 \in N$ and $a \in M$ the complex induced on a, b_1, b_2, u_p, v has facets $\{v, a, u_p\}, \{v, b_1, b_2\}, \{u_p, b_1, b_2\}$. This is J_1^*, and so $\mathcal{C}$ is not β-avoiding. □

Proposition 6.5. *Let $\mathcal{C}$ be the simplicial complex on q vertices with a vertex v such that $\text{link}_v(\mathcal{C}) = D_{m,n}$ with $m, n \geq 1$ and $(\mathcal{C} \setminus v)^* = G^{m+n-m'-n'}(\Delta_{m'} \sqcup \Delta_n)$ with $-1 \leq m' < m$ and $0 \leq n' \leq n$. Unless $m' = -1$ and $n' = n$, $\mathcal{C}$ has a proper minor that is not nuclear.*

Proof. Let M, N denote the sets of vertices such that in $(\text{link}_v(\mathcal{C}))^*$, the complex induced on M is Δ_m and the complex induced on N is Δ_n. Then some vertex $x \in N$ is a non-ghost in $(\mathcal{C} \setminus v)^*$. We show that $\text{link}_x(\mathcal{C})$ is not nuclear.

First we describe the facets of $\text{link}_x(\mathcal{C})$. We claim that the facets of $\text{link}_x(\mathcal{C})$ are precisely the sets containing $N-1-2$ vertices, where the omitted vertex pair is one of the following

(1) some $a \in M$ and some $b \in N \setminus \{x\}$ or
(2) v and some vertex g that is a ghost in $(\mathcal{C} \setminus v)^*$.

We now prove the claim. The facets of $\text{link}_x(\mathcal{C})$ that contain v are the sets whose complements are pairs of the first type since $\text{link}_v(\mathcal{C}) = D_{m,n}$. The facets of $\text{link}_x(\mathcal{C})$ that do not contain v are the sets whose complements are minimal non-faces of $(\mathcal{C} \setminus v)^*$ that do not become faces of $\text{link}_x(\mathcal{C})$ upon adding v. These minimal non-faces of $(\mathcal{C} \setminus v)^*$ are the ghost vertices. Therefore the facets of $\text{link}_x(\mathcal{C})$ that do not contain v are the sets of the second type.

We now show that $\text{link}_x(\mathcal{C})$ is not nuclear. The complex $\text{link}_x(\mathcal{C})$ cannot be $\Lambda(\mathcal{D})$ for any complex $\mathcal{D}$ since no facet lacks fewer than two vertices. Nor can $\text{link}_x(\mathcal{C})$ have any ghost vertices - $m \geq 1$ implies $\#M \geq 2$ and so every vertex in M is in a facet of the first type, every vertex of N is in a facet of the second type, and v is in every facet of the first type. It is clear that no vertex is in every facet, so $\text{link}_x(\mathcal{C})$ cannot be $\text{cone}^p(D_{k,l})$ nor $\text{cone}^p(\Delta_k \sqcup \Delta_l)$ for any $p \geq 1$. It is clear that $\text{link}_x(\mathcal{C})$ has more than 2 facets, so $\text{link}_x(\mathcal{C})$ cannot be $\Delta_k \sqcup \Delta_l$, nor Δ_k.

The only case left to check is $\text{link}_x(\mathcal{C}) = D_{k,l}$. If this is the case then we can partition the vertices of $\text{link}_x(\mathcal{C})$ into disjoint sets C_1, C_2 such that the facets of $\text{link}_x(\mathcal{C})$ are precisely the subsets of $C_1 \cup C_2$ that omit exactly one from each C_1 and C_2.

Assume $n' = n$ and $m' \geq 0$. Let $g \in M$ be a ghost vertex and without loss of generality assume $g \in C_1$. Facets of the first type imply $M \subseteq C_1$. Facets of the second type imply $v \in C_2$. Since $m' \geq 0$, there exists some $a \in M$ that is not a ghost in $(\mathcal{C} \setminus v)^*$. Snd so $(C_1 \cup C_2) \setminus \{a, v\}$ is a facet of $\text{link}_x(\mathcal{C})$. But this does not fit our earlier description of the facets of $\text{link}_x(\mathcal{C})$.

Now assume $-1 \leq m' < m$ and $0 \leq n' < n$. Then there exist $g' \in M$ and $g'' \in N$ that are ghosts in $(\mathcal{C} \setminus v)^*$. But this contradicts our earlier description of the facets of $\text{link}_x(\mathcal{C})$ - facets of the first kind imply that g', g'' are in different C_is and facets of the second kind imply that they are in the same C_i. □

Proposition 6.6. *Let $\mathcal{C}$ be a simplicial complex on q vertices. Let v be a vertex in $\mathcal{C}$ such that $\text{link}_v(\mathcal{C}) = D_{m,n}$ with $m, n \geq 1$. Assume $(\mathcal{C} \setminus v)^* = G^{m+1}(\Delta_n)$. Then $\mathcal{C} = D_{m,n+1}$.*

Proof. Let M, N denote the sets of vertices such that in $(\text{link}_v(\mathcal{C}))^*$, the complex induced on M is Δ_m and the complex induced on N is Δ_n. Then all the vertices of M and none

of the vertices of N are ghosts in $(\mathcal{C} \setminus v)^*$. We claim that each facet of $\mathcal{C}$ contains $N-2$ vertices, and the omitted pair is one of the following

(1) some $a \in M$ and some $b \in N$
(2) v and some $a \in M$.

To see this, note that if a facet contains v, then it must be of the first form since $\text{link}_v(\mathcal{C}) = D_{m,n}$. If a facet does not contain v, then its complement must be v, along with a minimal non-face of $(\mathcal{C} \setminus v)^*$. These minimal non faces are precisely the vertices in M. So $\mathcal{C} = D_{m,n+1}$ where M and $N \cup \{v\}$ are the sets of vertices for the Δ_m and Δ_{n+1} in $\mathcal{C}^*$ respectively. □

Proposition 6.7. *Let $\mathcal{C}$ be a simplicial complex on q vertices. Let v be a vertex in $\mathcal{C}$ such that $\text{link}_v(\mathcal{C}) = D_{m,n}$ with $m, n \geq 1$ and has no ghost vertices. Let M, N denote the sets of vertices such that in $(\text{link}_v(\mathcal{C}))^*$, the complex induced on M is Δ_m and the complex induced on N is Δ_n. Assume $(\mathcal{C} \setminus v)^* = G^{m+1}(\mathcal{D})$ where $\mathcal{D}$ is a nuclear complex on vertex set N with more than one facet. Then $\mathcal{C}$ has a proper minor that is not nuclear.*

Proof. Let $x \in M$. We show that $\text{link}_x(\mathcal{C})$ is not nuclear. We start by showing it is not Lawrence type. Let F be a facet of $\text{link}_x(\mathcal{C})$ that does not contain v. Then $F \cup \{x\}$ is a facet of $\mathcal{C} \setminus v$ so it must also lack some $a \in M \setminus \{x\}$. If F is a facet of $\text{link}_x(\mathcal{C})$ that contains v, then $F \cup \{x\} \setminus \{v\}$ is a facet of $\text{link}_v(\mathcal{C})$ and therefore lacks some $a \in M$ and $b \in N$. So $\text{link}_x(\mathcal{C})$ is not $\Lambda(\mathcal{F})$ for any complex $\mathcal{F}$.

We can see that $\text{link}_x(\mathcal{C})$ has more than two facets - since $m, n \geq 1$, there are at least two that include v and since $(C \setminus v)^* = G^{m+1}(\mathcal{D})$, there is at least one that does not. $\text{link}_x(\mathcal{C})$ cannot be $\text{cone}^p(\Delta_k \sqcup \Delta_l)$ for any $p \geq 0$.

All of the vertices of M are ghost vertices in $(\mathcal{C} \setminus v)^*$, and $(\mathcal{C} \setminus v)^*$ has a minimal non-face $S \subseteq N$ with at least two vertices. This implies that $\mathcal{C}$ contains a facet F such that $M \subseteq F$, there exist $b_1, b_2 \in N$ such that $b_1, b_2 \notin F$ and $v \notin F$ (since $\text{link}_v(\mathcal{C}) = D_{m,n}$). Facets in $\text{link}_x(\mathcal{C})$ that contain v have $N-1-2$ vertices - they lack some $a \in M \setminus \{x\}$ and some $b \in N$. Furthermore, $F \setminus \{x\}$ is a facet of $\text{link}_x(\mathcal{C})$ containing at most $N-1-3$ vertices. This means that $\text{link}_x(\mathcal{C})$ is not pure, and therefore not $\text{cone}^p(D_{k,l})$ for any $p \geq 0$. So $\text{link}_x(\mathcal{C})$ is a proper minor of $\mathcal{C}$ that is not nuclear. □

Lemma 6.8. *Let $\mathcal{C}$ be a β-avoiding simplicial complex on q vertices. Assume all proper minors of $\mathcal{C}$ are nuclear. If there exists a vertex v of $\mathcal{C}$ such that $\text{link}_v(\mathcal{C}) = \text{cone}^p(D_{m,n})$ for $p \geq 0$ and $m, n \geq 1$, then $\mathcal{C}$ is nuclear.*

Proof. We split this into two cases. For the first case, assume $p \geq 1$. Here, Propositions 3.9 and 3.18 give $\mathcal{C}^* \setminus v = \text{cone}^p(\Delta_m \sqcup \Delta_n)$. By Proposition 4.3, $\mathcal{C}^*$ is β-avoiding so in $\mathcal{C}^*$, v must connect to one of the simplices in the $\Delta_m \sqcup \Delta_n \subset \mathcal{C}^*$ to avoid inducing an independent set on three vertices. Now let u be a cone vertex. If v connects to only one such simplex, then v must also connect to u to avoid inducing P_4. If v connects to both such simplices, and v does not connect to u, then $\mathcal{C}^*$ has C_4 induced. In this case Proposition 4.4 implies $\mathcal{C}^* = \text{cone}^k(C_4)$ and so $\mathcal{C} = \text{cone}^k(\Delta_1 \sqcup \Delta_1)$. So we can assume v connects to at least one simplex, and to all cone vertices in $\mathcal{C}^* \setminus v$. From this we can see that $\mathcal{C}^*$ has no induced C_4. Since every proper minor of $\mathcal{C}^*$ is nuclear, Proposition 6.4

implies $\mathcal{C}^* = \text{cone}^{p+1}(\Delta_m \sqcup \Delta_n)$ or $\mathcal{C}^* = \text{cone}^p(\Delta_{m+1} \sqcup \Delta_n)$. Proposition 6.2 then implies that $\mathcal{C}$ is nuclear.

For the second case, assume $p = 0$. So $\text{link}_v(\mathcal{C}) = D_{m,n}$ with $m, n \geq 1$. Then $D_{m,n} \subseteq \mathcal{C} \setminus v$ and so $(\mathcal{C} \setminus v)^* \subseteq \Delta_m \sqcup \Delta_n$. Let M, N denote the sets of vertices of $\mathcal{C}$ such that in $(\text{link}_v(\mathcal{C}))^*$, the vertices of the Δ_m are M, and the vertices of the Δ_n are N. The non-ghost vertices of $(\mathcal{C} \setminus v)^*$ must form a nuclear complex, and the only disconnected nuclear complexes are of the form $\Delta_s \sqcup \Delta_t$. From this it follows that $(\mathcal{C} \setminus v)^*$ must be one of the following forms (without loss of generality)

(1) $\Delta_m \sqcup \Delta_n$
(2) $G^{m+n-m'-n'}(\Delta_{m'} \sqcup \Delta_{n'})$ with $-1 \leq m' < m$ and $0 \leq n' \leq n$, but not both $m' = -1$ and $n' = n$
(3) $G^{m+1}(\Delta_n)$
(4) $G^{m+1}(\mathcal{D})$ where $\mathcal{D}$ is a nuclear complex on N with more than one facet
(5) $G^{m+n+2}(\{\emptyset\})$
(6) $G^{m+n+2}(\{\})$.

We handle each possibility separately.

Case 1. Assume $(\mathcal{C} \setminus v)^* = \Delta_m \sqcup \Delta_n$. Then $(\mathcal{C} \setminus v)^*$ has no ghost vertices, and so $\mathcal{C} \setminus v$ is not of Lawrence type. So, by Lemma 4.7, $\mathcal{C} \setminus v = D_{m,n}$ and so $\mathcal{C} = \text{cone}^1(D_{m,n})$ with v as the cone point.

Case 2. Assume $(\mathcal{C} \setminus v)^* = G^{m+n-m'-n'}(\Delta_{m'} \sqcup \Delta_{n'})$ with $-1 \leq m' < m$ and $0 \leq n' \leq n$ but not both $m' = -1$ and $n' = n$. By Proposition 6.5, $\mathcal{C}$ has a proper minor M that is not nuclear. By induction, M is not β-avoiding and so neither is $\mathcal{C}$.

Case 3. Assume $(\mathcal{C} \setminus v)^* = G^{m+1}(\Delta_n)$. Then Proposition 6.6 implies that $\mathcal{C} = D_{m,n+1}$.

Case 4. Assume $(\mathcal{C} \setminus v)^* = G^{m+1}(\mathcal{D})$ where $\mathcal{D}$ is a nuclear complex on N with more than one facet. By Proposition 6.7, $\mathcal{C}$ has a proper minor M that is not nuclear. By induction, M is not β-avoiding and so neither is $\mathcal{C}$.

Case 5. Assume $(\mathcal{C} \setminus v)^* = G^{m+n+2}(\{\emptyset\})$. Then $\mathcal{C} \setminus v = \partial\Delta_{m+n+1}$. But $\text{link}_v\mathcal{C} = D_{m,n}$ and so v is not in any $m + n - 1$ simplex with the vertices of M and N. By Proposition 4.6, $\mathcal{C}$ is not β-avoiding.

Case 6. Assume $(\mathcal{C} \setminus v)^* = G^{m+n+2}(\{\})$. Then $\mathcal{C} \setminus v = \Delta_{m+n+1}$ and so $\mathcal{C} = \Lambda(D_{m,n})$ with v as the added vertex. □

Now we have all the necessary tools to prove Theorem 6.3.

Proof of Theorem 6.3. By Proposition 3.20, (1) implies (2). Each nuclear complex can be obtained by applying the unimodularity-preserving operations from Section 3 to $\Delta_n \sqcup \Delta_m$, which is unimodular by Proposition 3.21. Therefore (3) implies (1). We now show (2) implies (3).

Let $\mathcal{C}$ be a β-avoiding complex on q vertices. We show that $\mathcal{C}$ is nuclear by induction on q. For the base case, note that all simplicial complexes on 2 or fewer vertices are both β-avoiding and nuclear. We may assume that $\mathcal{C}$ has no ghost vertices since adding and removing ghost vertices does not affect the properties of being nuclear nor β-avoiding. In light of Propositions 4.4 and 5.4 and Lemma 5.3, we only need to consider the cases where the 1-skeleton of $\mathcal{C}$ is K_q, or the union of two complete graphs, glued along a common

nonempty clique. Therefore, we can choose a vertex v of $\mathcal{C}$ that is in a facet with every other vertex. Since $\mathcal{C}$ is β-avoiding, so is $\text{link}_v(\mathcal{C})$. By induction, $\text{link}_v(\mathcal{C})$ is also nuclear. There are five main cases.

Case 1. This is the case where $\text{link}_v(\mathcal{C}) = \Lambda(\mathcal{D})$ for a nuclear $\mathcal{D}$. Let F denote the big facet of $\mathcal{C}$ and let u be the vertex of $\mathcal{C}$ that is not in F. Then $\mathcal{C} = \Lambda(\mathcal{C} \setminus u)$, and $\mathcal{C} \setminus u$ is nuclear by induction.

Case 2. This is the case where $\text{link}_v(\mathcal{C})$ has ghost vertices. Since v connects to all other vertices, this is impossible.

Case 3. This is the case where $\text{link}_v(\mathcal{C}) = \text{cone}^p(\Delta_m \sqcup \Delta_n)$ with $p, m, n \geq 0$. We split this into the sub cases, the first where $m, n \geq 1$ and the second where $m = 0$.

Case 3.1 This is the sub case where $m, n \geq 1$. Proposition 3.9 gives $\mathcal{C}^* \setminus v = \text{cone}^p(D_{m,n})$ and so $\mathcal{C}^*$ has $D_{m,n}$ induced. Proposition 4.5 then implies that $\mathcal{C}^*$ is either $\text{cone}^{p+1}(D_{m,n})$, or $\text{cone}^p(D_{m,n})$ with v as a ghost vertex. In the first case, $\mathcal{C} = \text{cone}^{p+1}(\Delta_m \sqcup \Delta_n)$. In the second case, $\mathcal{C} = \Lambda(\text{cone}^p(\Delta_m \sqcup \Delta_n))$.

Case 3.2 This is the sub case where $m = 0$. Proposition 3.9 gives $\mathcal{C}^* \setminus v = \text{cone}^p(D_{0,n})$ which has a ghost vertex u. Then $\mathcal{C}^* \setminus u$ is nuclear by induction and since u is still a ghost vertex in $\mathcal{C}^*$, this implies $\mathcal{C}^*$ is nuclear. Proposition 6.2 implies that $\mathcal{C}$ is nuclear.

Case 4. This is the case where $\text{link}_v(\mathcal{C}) = \text{cone}^p(D_{m,n})$ with $p \geq 0$ and $m, n \geq 1$. By induction, all proper minors of $\mathcal{C}$ are nuclear. Then Lemma 6.8 implies that $\mathcal{C}$ is nuclear.

Case 5. This is the case where $\text{link}_v(\mathcal{C}) = \Delta_k$ with $k \geq -2$. Since $\text{link}_v(\mathcal{C})$ has no ghost vertices $k = q - 2$. So $\mathcal{C} = \Delta_{q-1}$. □

7. Unimodularity of Non-Binary Hierarchical Models

In this section we consider the unimodularity of hierarchical models that are non-binary; i.e. determining when $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ is unimodular for arbitrary $\mathbf{d}$. We begin by showing how Theorem 6.3 makes this problem more tractable. Unlike in the binary case, the Alexander dual does not always preserve unimodularity. For example, $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ is unimodular when $\mathcal{C} = \Delta_1 \sqcup \Delta_1$ and $d = (2, 2, 2, 3)$. However, $\mathcal{A}_{\mathcal{C}^*,\mathbf{d}}$ is not (see Proposition 7.9). Because of this, our proof of Proposition 3.10 is not valid in the non-binary case. The proposition statement does however generalize as we show with Corollary 7.4. Proposition 7.9 gives a list of known minimal pairs $(\mathcal{C}, \mathbf{d})$ with $\mathcal{C}$ nuclear where $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ is not unimodular. We then use this list to pose a simple question whose answer would be a step in the general classification.

Proposition 7.1. *Assume $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ is unimodular. Then for all $\mathbf{d}'$ such that $\mathbf{d}' \leq \mathbf{d}$ componentwise, $\mathcal{A}_{\mathcal{C},\mathbf{d}'}$ is unimodular.*

Proof. Note that $\mathcal{A}_{\mathcal{C},\mathbf{d}'}$ can be realized as a subset of columns of $\mathcal{A}_{\mathcal{C},\mathbf{d}}$. □

The special case where $\mathbf{d}' = \mathbf{2}$ gives us the following useful corollary.

Corollary 7.2. *If $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ is unimodular, then $\mathcal{C}$ is nuclear.*

So in order to classify all unimodular hierarchical models we only need to consider nuclear complexes. Therefore we can approach the general classification problem by looking at each nuclear $\mathcal{C}$ and identifying the minimal values of $\mathbf{d}$ that give rise to non-unimodular

$\mathcal{A}_{\mathcal{C},\mathbf{d}}$. Before proceeding with this, we show that the class of unimodular $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ is still closed under taking minors of $\mathcal{C}$. However Corollary 3.7 fails in the non-binary case; we cannot freely take the Alexander dual of $\mathcal{C}$. Because of this, our proof of Corollary 3.10 is not valid in the non-binary case. We will give an alternate proof that the class of unimodular $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ is closed under taking links in $\mathcal{C}$. We start with a useful proposition.

Proposition 7.3. *Let $A \in \mathbb{R}^{r\times n}$ be a unimodular matrix with columns $\{\mathbf{a}_i\}_{i=1}^n$. Assume $\mathbf{a}_n$ is nonzero. Let A' be the matrix that results when we project A onto the hyperplane orthogonal to $\mathbf{a}_n$. Then A' is unimodular.*

Proof. We may assume A has rank r since otherwise we can delete unnecessary rows. Let $B \in \mathbb{R}^{n\times(n-r)}$ have rank $n-r$ such that $AB^T = 0$. Denote the columns of B as $\{\mathbf{b}_i\}_{i=1}^n$. By Proposition 3.5, B is also unimodular. Denote the columns of A' as $\{\mathbf{a}'_i\}_{i=1}^{n-1}$. Then

$$\mathbf{a}'_i = \mathbf{a}_i - \frac{\langle \mathbf{a}_i, \mathbf{a}_n\rangle}{\|\mathbf{a}_n\|^2}\mathbf{a}_n.$$

Let B' be the matrix with columns $\{b_i\}_{i=1}^{n-1}$. Then B' has rank $n-r$ or $n-r-1$. The second case implies that $\mathbf{b}_n$ is a coloop in the matroid underlying B. Since the matroids underlying A and B are duals, this would imply that $\mathbf{a}_n = 0$. But then $A' = A$ which we assumed to be unimodular. So we can assume that $\operatorname{rank}(B') = \operatorname{rank}(B) = n-r$ and that $\mathbf{a}_n \neq 0$. Therefore $\operatorname{rank}(A') = r-1$ and so the dimension of its kernel is $n-r$, which is the rank of B'. We claim that $A'(B')^T = 0$. From this it follows by Proposition 3.5 that A' is unimodular.

Now we prove the claim. Let $\mathbf{b}_i$ be a column of B' (so $1 \le i \le n-1$). Letting b_{ji} denote the jth entry of $\mathbf{b}_i$, we have

$$A'\mathbf{b}_i^T = \sum_{j=1}^{n-1} b_{ji}\left(\mathbf{a_j} - \frac{\langle \mathbf{a}_i, \mathbf{a}_n\rangle}{\|\mathbf{a}_n\|^2}\mathbf{a}_n\right).$$

Note that $b_{jn}\left(\mathbf{a}_n - \frac{\langle \mathbf{a}_n,\mathbf{a}_n\rangle}{\|\mathbf{a}_n\|^2}\mathbf{a}_n\right) = 0$, so we can add the nth term to the above sum. This enables us to break it up as follows

$$\sum_{j=1}^{n} b_{ji}\mathbf{a}_j - \frac{1}{\|\mathbf{a}_n\|^2}\sum_{j=1}^{n} b_{ji}\langle \mathbf{a}_j, \mathbf{a}_n\rangle\mathbf{a}_n.$$

The first term is $A\mathbf{b}_i^T = 0$. The second term is $\left(\frac{1}{\|\mathbf{a}_n\|^2}\mathbf{a}_n^T A\mathbf{b}_i^T\right)\cdot \mathbf{a}_n = 0$. So the claim is proven. □

Corollary 7.4. *Assume $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ is unimodular. Let v be a vertex of $\mathcal{C}$ and let $\mathbf{d}'$ denote the vector obtained by deleting the entry for v from $\mathbf{d}$. Then $\mathcal{A}_{\operatorname{link}_v\mathcal{C},\mathbf{d}'}$ is unimodular.*

Proof. Let $A = \mathcal{A}_{\mathrm{link}_v \mathcal{C}, \mathbf{d}'}$ and let $B = \mathcal{A}_{\mathcal{C}\setminus v, \mathbf{d}'}$. By the remark following Lemma 2.2 in [7] and the lemma itself, we can write

$$\mathcal{A}_{\mathcal{C},\mathbf{d}} = \begin{bmatrix} A & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & A & \dots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \dots & A \\ B & B & \dots & B \end{bmatrix}.$$

Assume $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{l \times n}$ and so $\mathcal{A}_{\mathcal{C},\mathbf{d}} \in \mathbb{R}^{dm+l, dn}$. By Proposition 2.9(3), we may remove rows from $\mathcal{A}$ to make $A, B, \mathcal{A}$ all have full row rank. So assume that they do. Let $\mathcal{A}'$ denote the matrix that results when we project $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ onto the subspace orthogonal to the last $(d-1)n$ columns. Then if $1 \leq i \leq n$, the ith column of $\mathcal{A}'$ can be expressed as the ith column of $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ minus a linear combination of the last $(d-1)n$ columns of $\mathcal{A}_{\mathcal{C},\mathbf{d}}$, all of which are 0 in the top m rows. So this means that the top m rows of $\mathcal{A}'$ are

$$\begin{pmatrix} A & \mathbf{0} & \dots & \mathbf{0} \end{pmatrix}.$$

Furthermore, since A and $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ both have full row rank, the final $(d-1)n$ columns of $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ have rank $(d-1)m + l$. Therefore $\mathcal{A}'$ has rank m. Since A also has rank m, we may delete the bottom $(d-1)m + l$ rows of $\mathcal{A}'$ without affecting the rowspace and therefore unimodularity. So the matrix $\begin{pmatrix} A & \mathbf{0} & \dots & \mathbf{0} \end{pmatrix}$, and therefore A, is unimodular. □

Our proof of Proposition 3.1 is still valid in the non-binary case. Therefore, we have the following.

Proposition 7.5. *Assume $\mathcal{C}'$ is a minor of $\mathcal{C}$. If $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ is unimodular, then so is $\mathcal{A}_{\mathcal{C}',\mathbf{d}'}$ where $\mathbf{d}'$ is the restriction of $\mathbf{d}$ to the vertices that are in $\mathcal{C}'$.*

The proofs of Propositions 3.12, 3.17 generalize to the non-binary setting and so we have the following.

Proposition 7.6. *If the pair $(\mathcal{C}, \mathbf{d})$ gives rise to unimodular $\mathcal{A}_{\mathcal{C},\mathbf{d}}$, so do $(\mathrm{cone}^p(\mathcal{C}), \mathbf{d}')$ and $(G(\mathcal{C}), \mathbf{d}'')$ where $\mathbf{d}' = \begin{pmatrix} \mathbf{d} & c_1 & \dots & c_p \end{pmatrix}$ and $\mathbf{d}'' = \begin{pmatrix} \mathbf{d} & c \end{pmatrix}$ for any $c, c_1, \dots, c_p \geq 2$.*

Proof. In the case of a cone, the matrix $\mathcal{A}_{\mathrm{cone}^p(\mathcal{C}),\mathbf{d}'}$ is a block diagonal matrix:

$$\mathcal{A}_{\mathrm{cone}^p(\mathcal{C}),\mathbf{d}'} = \begin{pmatrix} \mathcal{A}_{\mathcal{C},\mathbf{d}} & 0 & \cdots & 0 \\ 0 & \mathcal{A}_{\mathcal{C},\mathbf{d}} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \mathcal{A}_{\mathcal{C},\mathbf{d}} \end{pmatrix},$$

with $c_1 c_2 \cdots c_p$ blocks down the diagonal. In the case of adding a ghost vertex, the matrix $\mathcal{A}_{G(\mathcal{C}),\mathbf{d}'}$ has repeated columns:

$$\mathcal{A}_{G(\mathcal{C}),\mathbf{d}''} = \begin{pmatrix} \mathcal{A}_{\mathcal{C},\mathbf{d}} & \mathcal{A}_{\mathcal{C},\mathbf{d}} & \cdots & \mathcal{A}_{\mathcal{C},\mathbf{d}} \end{pmatrix}$$

with c blocks. □

We can also extend Proposition 3.21.

Proposition 7.7. *Let $\mathcal{C} = \Delta_m \sqcup \Delta_n$ and $\mathbf{d} \in \mathbb{Z}_{\geq 2}^{m+n+2}$. Let M, N denote the vertex sets of $\mathcal{C}$ in the Δ_m, Δ_n respectively. Let $\mathcal{D}$ be the complex with facets $\{1,2\}$ and let $\mathbf{e} = (e_1, e_2)$ where*

$$e_1 = \prod_{v \in M} d_v \qquad e_2 = \prod_{v \in N} d_v.$$

Then $\mathcal{A}_{\mathcal{C},\mathbf{d}} = \mathcal{A}_{\mathcal{D},\mathbf{e}}$. This matrix is unimodular.

Proof. We can see that $\mathcal{A}_{\mathcal{C},\mathbf{d}} = \mathcal{A}_{\mathcal{D},\mathbf{e}}$ by inspection. The matrix $\mathcal{A}_{\mathcal{D},(e_1,e_2)}$ is the vertex edge incidence matrix of a complete bipartite graph with e_1 and e_2 vertices in each set of the partition. Such vertex-edge incidence matrices are examples of network matrices and are hence totally unimodular [12, Ch. 19]. □

We also note that Proposition 3.15 holds in a slightly more general setting.

Proposition 7.8. *Let $(\mathcal{C}, \mathbf{d})$ be such that $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ is unimodular. Then $\mathcal{A}_{\Lambda(\mathcal{C}),\mathbf{d}'}$ is also unimodular if $\mathbf{d}' = \begin{pmatrix} \mathbf{d} & 2 \end{pmatrix}$.*

Proof. The proof here is similiar to the proof for Proposition 3.15. Note that $\Lambda(\mathcal{A}_{\mathcal{C},\mathbf{d}}) = \mathcal{A}_{\Lambda(\mathcal{C}),\mathbf{d}'}$. Hence Theorem 7.1 in [14] implies the proposition. □

Propositions 7.1 and 7.5 suggest that we might be able to use a short list of pairs $(\mathcal{C}, \mathbf{d})$ that give rise to non-unimodular $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ to eliminate the remaining non-unimodular pairs. The following proposition lists all non-binary minimally non-unimodular complexes that we presently know.

Proposition 7.9. *The following pairs give rise to non-unimodular $\mathcal{A}_{\mathcal{C},\mathbf{d}}$*

(1) *$\mathcal{C} = \Lambda(\Delta_0 \sqcup \Delta_0)$ with facets $\{12, 23, 13\}$ and $\mathbf{d} = (3,3,3)$*
(2) *$\mathcal{C} = D_{1,1}$ with facets $\{12, 23, 34, 14\}$ and $\mathbf{d} = (2,2,2,3)$*
(3) *$\mathcal{C} = \Lambda(G(\Delta_0 \sqcup \Delta_0))$ with facets $\{12, 13, 234\}$ and $\mathbf{d} = (4,2,2,2)$*
(4) *$\mathcal{C} = \Lambda(D_{1,1})$ with facets $\{1234, 125, 235, 345, 145\}$ and $\mathbf{d} = (2,2,2,2,3)$.*

Proof. By Table 1 in [9], (1) is not unimodular. Computing a Graver basis with 4ti2 [1] shows that (2) and (4) are not unimodular. Computations are available on our website [3]. If $\mathcal{C}$ and $\mathbf{d}$ are as in (3) then $\mathcal{A}_{\mathcal{C},\mathbf{d}} = \mathcal{A}_{J_1^*,\mathbf{2}}$ which is not unimodular by Proposition 4.1. □

From these non-unimodular examples and the constructive results earlier in the section, we can classify all unimodular hierarchical models that are nuclear with nucleus $D_{m,n}$, $m, n \geq 1$.

Theorem 7.10. *Let $\mathcal{C}$ be a nuclear complex with nucleus $D_{m,n}$ with $m, n \geq 1$. Then $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ is unimodular if and only if*

(1) *$d_v = 2$ for all v in the original $D_{m,n}$ and*
(2) *$d_v = 2$ for all v corresponding to a Lawrence lifting.*

Proof. The fact that all such $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ are unimodular follows from applying Propositions 7.6 and 7.8. We now show that these are the only unimodular $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ with this type of complex $\mathcal{C}$.

Note that since taking cone vertices commutes with adding ghost vertices and Lawrence liftings and this always preserves unimodularity, we can assume there are no cone vertices. Now if $\mathcal{C}$ is a nuclear complex obtained by successively adding ghost vertices and Lawrence vertices to $D_{m,n}$, then the induced subcomplex on just the $D_{m,n}$ and Lawrence vertices is an iterated Lawrence lifting of $D_{m,n}$. If any of the vertices of v that are Lawrence vertices have $d_v > 2$, then by taking suitable links, one obtains the complex isomorphic to item (4) in Proposition 7.9. On the other hand, if $d_v > 2$ for some v in the original $D_{m,n}$, then by taking suitable links, one obtains the complex isomorphic to item (2) in Proposition 7.9. In summary, if the conditions of Theorem 7.10 are not satisfied, there is a minor of $\mathcal{C}$ that is not unimodular. □

It remains to consider the unimodularity of $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ when $\mathcal{C}$ is a nuclear complex with nucleus $\Delta_m \sqcup \Delta_n$. We do not have a general characterization in this case. By Propositions 7.6 and 3.18, we may assume that $\mathcal{C}$ has no cone vertices. By Proposition 7.7, we may assume that $m, n = 0$. Note that $\Lambda(\Delta_0 \sqcup \Delta_0) = \partial\Delta_2$, and our results above (or [9]) show that this is unimodular if and only if some $d_v = 2$. Similarly, the iterated Lawrence liftings $\Lambda^p(\Delta_0 \sqcup \Delta_0) = \partial\Delta_{p+1}$ is unimodular if and only if at most two of the d_v are greater than 2.

We begin moving into uncharted territory when we consider cases where ghost vertices are involved. The simplest case to consider is the complex $\Lambda(G(\Delta_0 \sqcup \Delta_0))$ which is the complex on $\{1, 2, 3, 4\}$ with facets $\{12, 13, 234\}$. Here 1 is the Lawrence vertex and 4 is the ghost vertex. By Propositions 7.5 and 7.9 (3), we must have $d_1 \leq 3$. If $d_1 = 2$ then Proposition 7.8 implies that $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ will be unimodular. So we only need to consider the case where $d_1 = 3$. Propositions 7.1, 7.5, and 7.9 (1) imply that one of d_2 or d_3 is 2, so assume $d_2 = 2$. This leaves us with the following question, which would classify unimodularity for $\mathcal{C} = \Lambda(G(\Delta_0 \sqcup \Delta_0))$.

Question 7.11. Let $\mathcal{C} = \Lambda(G(\Delta_0 \sqcup \Delta_0))$ be the complex on $\{1, 2, 3, 4\}$ with facets $\{12, 13, 234\}$. Let $\mathbf{d} = (3, 2, d_3, d_4)$. For what values of d_3, d_4 is the matrix $\mathcal{A}_{\mathcal{C},\mathbf{d}}$ unimodular?

Acknowledgments

Daniel Irving Bernstein was partially supported by the US National Science Foundation (DMS 0954865). Seth Sullivant was partially supported by the David and Lucille Packard Foundation and the US National Science Foundation (DMS 0954865).

References

[1] 4ti2 team. 4ti2—a software package for algebraic, geometric and combinatorial problems on linear spaces. Available at www.4ti2.de.

[2] Daniel Irving Bernstein and Seth Sullivant. Normal binary hierarchical models. `ArXiv:1508.05461`, 2015.

[3] Daniel Irving Bernstein and Seth Sullivant. Supplementary materials. http://www4.ncsu.edu/~smsulli2/Pubs/UnimodularWebsite/Unimodular.html, 2016.

[4] Yuguo Chen, Ian H. Dinwoodie, and Seth Sullivant. Sequential importance sampling for multiway tables. *Ann. Statist.*, 34(1):523–545, 2006.

[5] Persi Diaconis and Bernd Sturmfels. Algebraic algorithms for sampling from conditional distributions. *Ann. Statist.*, 26(1):363–397, 1998.

[6] Serkan Hoşten and Seth Sullivant. Gröbner bases and polyhedral geometry of reducible and cyclic models. *J. Combin. Theory Ser. A*, 100(2):277–301, 2002.

[7] Serkan Hoşten and Seth Sullivant. A finiteness theorem for Markov bases of hierarchical models. *J. Combin. Theory Ser. A*, 114(2):311–321, 2007.

[8] Tomáš Kaiser. Minors of simplicial complexes. *Discrete Applied Mathematics*, 157(12):2597 – 2602, 2009. Second Workshop on Graph Classes, Optimization, and Width Parameters.

[9] Hidefumi Ohsugi and Takayuki Hibi. Non-very ample configurations arising from contingency tables. *Annals of the Institute of Statistical Mathematics*, 62(4):639–644, 2010.

[10] Ignacio Ojeda and Alberto Vigeneron-Tenorio. Indispensable binomials in semigroup ideals. *Proc. Amer. Math. Soc.*, 138:4205–4216, 2010.

[11] Johannes Rauh and Seth Sullivant. Lifting markov bases and higher codimension toric fiber products. `ArXiv:1404.6392`, 2014.

[12] Alexander Schrijver. *Theory of Linear and Integer Programming*. Wiley-Interscience Series in Discrete Mathematics. John Wiley & Sons, Ltd., Chichester, 1986. A Wiley-Interscience Publication.

[13] Paul Seymour. Decomposition of regular matroids. *J. Combin. Theory Ser. B*, 28(3):305–359, 1980.

[14] Bernd Sturmfels. *Gröbner Bases and Convex Polytopes*, volume 8 of *University Lecture Series*. American Mathematical Society, Providence, RI, 1996.

[15] Seth Sullivant. Small contingency tables with large gaps. *SIAM Journal of Discrete Mathematics*, 18(4):787–793, 2005.

[16] Seth Sullivant. Normal binary graph models. *Ann. Inst. Statist. Math.*, 62(4):717–726, 2010.

[17] W. T. Tutte. A homotopy theorem for matroids i,ii. *Trans. Amer. Math. Soc.*, 88:144–174, 1958.

Department of Mathematics, North Carolina State University, Raleigh, NC 27695
E-mail address: `dibernst@ncsu.edu`
E-mail address: `smsulli2@ncsu.edu`